

THE DYNAMICAL EVOLUTION OF NARROW LINE REGIONS

M. A. Dopita, G. V. Bicknell, R. S. Sutherland & C. J. Saxton

Research School of Astronomy & Astrophysics, The Australian National University.

RESUMEN

El resumen será traducido al español por los editores. We review recent progress by the RSAA team in elucidating the dynamical evolution of the various classes of Narrow Line Regions (NLR) in active galaxies.

ABSTRACT

We review recent progress by the RSAA team in elucidating the dynamical evolution of the various classes of Narrow Line Regions (NLR) in active galaxies.

Key Words: **ISM:HYDRODYNAMICS — ISM: JETS AND OUTFLOWS — GALAXIES:AGN — GALAXIES:SEYFERT — GALAXIES:EVOLUTION**

1. NLR ZOOLOGY

Optical spectroscopy and dynamical analysis has revealed that the so-called narrow line regions (NLR) are an almost ubiquitous signature of the presence of an active nucleus in a galaxy (AGN). The optical diagnostic diagrams of Baldwin, Phillips & Terlevich (1981), Veilleux & Osterbrock (1987) and Osterbrock et al. (1992) serve to classify a region as an NLR, and such regions can be further sub-divided into two general categories: Seyfert narrow line regions (NLRs) and the Low Excitation Nuclear Emission Line Regions (LINERs), which will not be discussed here

Homogenous sets of high-quality spectrophotometric data are now available for the NLR of most classes of AGN. Notable amongst these are complete sample of southern elliptical galaxies by Phillips et al. (1985), the similar-sized sample of nearby northern spirals by Ho, Filippenko & Sargent (1995), the compilations of Veilleux & Osterbrock (1987), and Véron & Véron-Cetty (2000), the Seyfert galaxy study of Veilleux (1991a,b,c), the compact radio-luminous sample of Gelderman & Whittle (1994), the luminous infrared galaxy survey of Kim et al. (1995), the ultraluminous infrared galaxy survey of Kim et al. (1998) and Veilleux, Kim & Sanders (1999), the warm IRAS sample of Kewley et al. (2001a), and the radio-excess IRAS galaxies (Drake et al, 2003, in press). NLR are found not only in nearby galaxies, but they also appear to be a ubiquitous feature of QSOs Francis et al. (1991), and they provide the dominant form of near-nuclear optical emission in the distant high-power radio sources (Best, Röttgering & Longair 2000a; Best, Röttgering & Longair 2000b; Inskip et al. 2002a; Inskip et al. 2002b; De Breuck 2000).

There is increasing evidence that the narrow-line regions (NLRs) associated with many classes of active galactic nuclei (AGN) have a complex dynamical and excitation evolution. Evidence for a cocoon of strong, auto-ionising and radiative shocks is particularly compelling for luminous classes of radio galaxies (see the theory by Dopita & Sutherland 1995; Dopita & Sutherland 1996 and Bicknell Dopita & O’Dea 1997). These include the steep-spectrum radio sources (CSS), Fanti et al. 1990, unbeamed Gigahertz-peaked sources (GPS) (see the recent review by O’Dea et al. 1998), compact symmetric objects (CSO) (Wilkinson et al. 1994) or compact double sources (CD) (Phillips & Mutel 1982). Together, these represent an appreciable fraction (10-30%) of luminous radio sources. Not only are such sources very luminous at radio frequencies, but they also are very luminous in optical emission lines, and spectra by Gelderman & Whittle (1994) reveal intense “narrow line” emission with line ratios similar to those of Seyfert 2 galaxies. For these objects, the line emission scales with radio power, and the continuity of properties across these different classes of sources argues strongly that the kinetic energy supplied by the radio-emitting jets may provide a substantial fraction of the power radiated at other wavelengths by shocked gas associated with the NLRs of these galaxies.

Evidence for excitation by both shock and photons has been adduced from the powerful high-redshift radio galaxies. The study by Best et al. (2000a,b) revealed that for the 3C radio galaxies ($z \sim 1$) radio lobes small, they are predominantly shock-excited, but when the radio lobes have burst out into intergalactic space the ionised gas left be-

hind is predominantly photoionized. Similar results were obtained by De Breuck (2000) for the very high redshift radio galaxies.

The Seyfert galaxies are generally found in spiral galaxy hosts, often display lines of high-excitation, and evidence of non-gravitational motions, but are radio-quiet. Dynamical signatures of strong shocks are apparent in only 5 – 10% of cases (eg. Whittle 1996). Many of these found amongst the more radio-luminous galaxies including Mrk78 (Pedlar et al. 1989), NGC2992 Allen et al. (1999) or Mrk 1066 Bower et al. (1995). The weak collimated radio-jets frequently seen in such objects often show close correlations between the radio and the optical morphology (Allen et al. 1999; Axon et al. 1998; Bower et al. 1995; Capetti et al. 1995; Falcke, Wilson & Simpson 1998; Haniff, Wilson & Ward 1988; Whittle et al. 1988). Power requirements are modest, typically $10^{41} - 10^{44}$ ergs s $^{-1}$ *c.f.* the luminous radio sources ($10^{45} - 10^{46}$ erg s $^{-1}$). The remainder of Seyfert galaxies appear to be photoionised (Evans et al. 1999).

The Infrared Astronomical Satellite (IRAS) revealed a large population of galaxies with intense circum-nuclear star formation and which emit the bulk of their radiation in the infrared. Many of these have $\log(L_{FIR}) > 11.0$ and are referred to as luminous infrared galaxies (LIRGs). Although it is clear that the IR luminosity derives from dust reprocessing of other sources of luminosity in the galaxy, the nature of the nuclear source is still in debate. Most of the ultraluminous sources (ULIRGS; eg Goldader et al 1995) and the majority of lower luminosity galaxies Kim et al. (1995) are star formation dominated. Indeed, Condon, Anderson & Helou (1991) concluded that the far infrared luminosity and radio properties of LIRGs can be explained entirely by compact nuclear starburst events. However, Sanders et al. (1988) have argued that LIRGs contain a dust enshrouded AGN and Vielleux, Sanders & Kim (1997) have used near IR and optical spectroscopy to search for broad emission lines, indicative of AGN, which they found in some 25-30% of ULIGs of their sample, and Lonsdale, Smith & Lonsdale (1993) concluded that Active Galactic Nuclei (AGN) are the dominant powering mechanism. It is most likely that both mechanisms contribute to the overall energy output, and where each provides a similar flux, the nuclear spectra are “composite”. Such spectra are particularly common in luminous IR galaxy samples. They have been identified by Kim et al. (1995), Kim et al. (1998) and Veilleux, Kim & Sanders (1999), the studies of Véron, Gonçalves,

& Véron-Cetty (1997), Gonçalves, Véron-Cetty & Véron (1999) and in the extensive survey of warm IR galaxies by Kewley et al. (2001a). They are also found in radio-excess IRAS galaxies, which are a rare but important class of radio-intermediate sources. Drake et al. (2002) has shown that 40-45% of such objects are compact and have steep radio spectral indices, *i.e.* they fit the definition of CSS or GPS radio sources.

From all of this, it should be clear that both shocks and photoionization from the central nucleus are important in exciting the NLR. In this review, we will attempt to provide observational and theoretical insight into the dynamical evolution of these various classes of NLR objects, and infer under what circumstances each of these excitation mechanisms may dominate.

2. MODELS OF THE DYNAMICAL EVOLUTION OF THE NLR

2.1. The GPS/CSS Sources

In their initial phases of activity the relativistic jets produced by AGN are likely to interact strongly with the interstellar medium (ISM). Thus, the early evolution of the lobe is likely to provide insight into both the nature of the circum-nuclear ISM and of the jet itself. Modulo issues relating to the absolute power of the radio jet, the youngest AGN are probably also the smallest, and therefore an appropriate place to start our examination of the dynamical evolution of AGN is the gigahertz-peak sources (GPS) and the compact symmetric sources (CSS), which are likely to be triggered by the merger of a low-mass system onto a massive and evolved Elliptical galaxy. In what follows we draw upon the results of Bicknell Saxton & Sutherland (2002).

In his early models of these sources Begelman 1996 conjectured that the mean pressure in the radio hot-spot of these sources is in constant ratio to the pressure of the cocoon of hot and relativistic plasma which surrounds the jet. If the ambient ISM declines in density as a power law of the distance from the nucleus, $n(R) = n_0 (R/R_0)^{-\delta}$, then Begelman’s conjecture is valid for $\delta \sim 2$. This conjecture has been verified in detail (Carvalho & O’Dea 2002a; 2002b). The model of Bicknell Dopita & O’Dea (1997), which allows for the expansion losses provides the velocity of advance, v_B , of the bow-shock into a uniform medium following such a power-law with radius:

$$\frac{v_B}{c} \approx 0.056 \left[\frac{F_{46}}{n_{0.01}} \right]^{1/3} \left[\frac{R}{\text{kpc}} \right]^{(\delta-2)/3} \quad (1)$$

where here $n_{0.01}$ is the density at $R = 1$ kpc in units of 0.01 cm^{-3} and the jet power F_{46} is given in terms of $10^{46} \text{ ergs s}^{-1}$.

The optical emission in the CSS sources shows clear signatures of shock excitation, with systematic velocity offsets of $300 - 500 \text{ km s}^{-1}$, lines broadened and split by $\sim 500 \text{ km s}^{-1}$, and a strong alignment between the radio jets and optical emission (de Vreis 1997, 1999). Whilst such effects were predicted by the BDO model, the shocks in this model were produced by radiative wall shocks, which would require relatively high ISM densities. This would permit only a low velocity of advance of the bow-shock consistent with an expansion velocity of less than 1000 km s^{-1} in the walls of the cocoon, which is inconsistent with recent observations which suggest instead 0.1-0.3 times the speed of light. Also, the observations of de Vreis (1999) also show that the line emission is not concentrated around the head of the radio lobe, but trails behind it in the walls.

A likely explanation is that the majority of the ISM is in fact characterized by a low density medium with a strongly radially declining density gradient, but with much denser clouds are embedded in it. This is entirely consistent with what we know about the structure of the ISM in galaxies, with a cold and/or molecular component embedded in a hot medium at coronal temperatures, which in elliptical galaxies has a temperature of order $(1-2) \times 10^7 \text{ K}$ and $\delta \sim 3/2$. Irwin & Sarazin (1996). In such a cloudy medium, the overpressure in the cocoon surrounding the jets drives slower, radiative shocks into the dense clouds. For clouds lying outside the jet core itself, the ratio of the velocity of advance of the jet and the cloud shock velocity, v_S is given in terms of the cloud density, and the intercloud density in its vicinity, by

$$\frac{v_B}{v_S} = \zeta^{1/2} \left(\frac{n_C}{n_{IC}} \right)^{1/2} \quad (2)$$

where $\zeta \sim 10 - 100$ is the ratio of the ram pressure of the jet and pressure in the cocoon of shocked gas surrounding it.

Cloud shocks will become radiative when the cooling timescale behind the shock, τ_{cool} , is comparable to the jet expansion dynamical timescale, τ_{dyn} . The cooling timescale is given by radiative shock models (Dopita & Sutherland 1995, Dopita & Sutherland 1996), and in the velocity range $200 - 900 \text{ km s}^{-1}$ can be approximated as:

$$\tau_{\text{cool}} \approx 23000 n_C^{-1} \left[\frac{v_S}{300 \text{ km s}^{-1}} \right]^{3.9} \text{ yrs.} \quad (3)$$

The dynamical timescale is:

$$\tau_{\text{dyn}} = 32600 \left[\frac{v_B}{0.1c} \right]^{-1} \left[\frac{R}{\text{kpc}} \right] \text{ yrs.} \quad (4)$$

Approximate equality of these timescales when combined with equation (2) means that the cloud to intercloud density contrast has to be at least 10^3 and possibly as large as 10^4 , consistent with cloud temperatures of $10^3 - 10^4 \text{ K}$, or turbulent motions of order 10 km s^{-1} .

Modelling such a large density contrast (including radiative cooling) within a hydrodynamic code is a challenging business. Sutherland, Bisset & Bicknell (2002) have developed a code able to tackle this based on the VH-1 code of the Virginia group (see <http://wonka.physics.ncsu.edu/pub/VH-1>). This (originally adiabatic) PPM code has undergone numerous changes in its Riemann solver, and it features an “oscillation filter” designed to suppress a stripping numerical instability seeded by numerical noise, driven by directional splitting and exacerbated by radiative cooling. This code has recently been ported to the MPI parallel environment.

Some results of the 2D implementation of this code are shown in Figure 1, which shows a pair of clouds being shocked by the passage of the jet near them. When clouds are submitted to an overpressure large enough to drive *non-radiative* shocks into them, the clouds are rapidly ablated and torn apart, which entrains matter into the cocoon as a whole. However, when the shocks are radiative, the clouds cool and are crushed to a density which tends to put them into pressure equilibrium with the surrounding jet or cocoon environment. This makes them much more resistant to destruction on short timescales, considerably extending the cloud destruction time scale. Simulations of radio-lobe/cloud interactions are available as movies on <http://macnab.anu.edu.au/radiojets/gps>.

The evolution of the jet itself depends critically on the filling factor of the clouds. With low filling factor, the jet is bent or diverted by interaction with clouds, but is not entirely disrupted. In these cases, the evolution is like the jittering “dentist’s drill” propagation envisioned by Sheuer (1982). However, above a certain critical filling factor, the jet becomes “frustrated”, and is broken up into a series of channels reminiscent of a river delta (see Figure 2). In this case, the evolution of the whole cocoon is more like a cloudy stellar wind bubble. We should emphasize that jet shown in figure 2 is highly supersonic; its Mach number is 130. However, the jet kinetic energy is rapidly thermalised so that the




Fig. 1. Shocks induced by the passage of the jet are seen sweeping through clouds embedded in the jet cocoon. The grayscale is reversed at low levels so as to show the jet. Material which has been ablated from the clouds adds to the pressure of the cocoon.

pressure in the cocoon is very high; $\sim M^2$ times the background pressure.

2.2. *Hi-Z Radio Sources*

The theory and models discussed above suggest that, in their early phases, the jet and circum-nuclear ISM interact very strongly, and that the early evolution of the NLR is characterized by the presence of strong radiative shocks. Under such circumstances the “auto-ionizing” fast shock model of Dopita & Sutherland (1995) and Dopita & Sutherland (1996) would provide the dominant source of excitation.

If this is true in the case of the GPS/CSS sources, it will apply even more strongly to the gas-rich strong radio sources which are found in abundance in the high-redshift universe. Such galaxies present us with a statistically significant sample in which to study UV line ratio behaviour and so investigate the fraction of the NLR emission produced by shocks, and the fraction by photoionization. The study by Best *et al.* (2000a,b) revealed an extraordinary result for powerful 3C radio galaxies with $z \sim 1$. They find that both the UV line profiles and the UV line ratio diagnostics imply that, when the scale of the radio lobes is such that they are still able to interact with the gas in the vicinity of the galaxy, they are predominantly shock-excited, but when the lobe has burst out into intergalactic space, the ionised gas left behind is pre-




Fig. 2. The later evolution of a 2-D jet surrounded by a set of dense clouds. Above a critical filling factor, the jet is disrupted and “percolates” through the channels which it has been able to open up.

dominantly photoionized. The ratio of fluxes in the different classes of source suggests that the energy flux in the UV radiation field is about 1/3 of the energy flux in the jets. Thus, both shocks and photoionization are important in the overall evolution of radio galaxies. This result, confirmed by Inskip *et al.* (2002), proves that the properties of the radio jet are intimately connected with the central engine.

Very distant radio galaxies have been recently studied by De Breuck (2000). He finds that diagnostic diagrams involving C IV, He II and C III] fit to the pure photoionization models, but that the observed C II]/C III] requires there to be a high-velocity shock present. He argues that composite models would be required to give a self-consistent description of all the line ratios, and that these may require a mix of different physical conditions as well.

Such sources are uniquely associated with massive gas-rich multi- L_* galaxies in the early universe ($< 2 - 3$ Gyr). They display a strong “alignment effect”, with regions of very high star formation rate ($> 1000 M_\odot \text{ yr}^{-1}$), and emission line gas having the spectral characteristics of the NLR extended along the direction of the steep-spectrum radio lobes. In this objects, the radio jet appears to be driving strong shocks into the galaxian ISM (evidenced by extensive Ly- α haloes; Reuland *et al.* 2003) which in turn triggers enormous star formation in the sur-

rounding cocoon. A fine example of such a source is provided by the $z \sim 3.8$ radio galaxy 4C 41.17 which has recently been studied in detail by Bicknell et al. (2000). This object consists of a powerful “double-double” radio source embedded in a 190×130 kpc Ly α halo Reuland et al. (2003) and shows strong evidence for jet-induced star formation at $3000 M_{\odot} \text{ yr}^{-1}$ associated with the inner radio jet. This is apparently induced by the strong dynamical interaction of the inner jet with the shocked and compressed gas in the wall of the cocoon created by the passage of the outer jet. Shock-induced starformation in jet walls was proposed in the context of Seyfert galaxies by Steffen et al. (1997). In 4C 41.17, the outer jet also appears to have induced a large-scale outflow with velocities in excess of 500 km s^{-1} in the line-emitting gaseous halo. Thus we may be seeing the “end of the beginning” in which the central super-massive black hole has finally become large enough to drive the whole accreting envelope of gas into outflow, triggering a last and spectacular burst of star formation in the process.

2.3. Seyferts

Seyferts are radio-quiet, and most appear to be dominated by photoionization by the central object. The dynamical signatures of strong shocks are apparent in only 5 – 10% of Seyferts (eg. Whittle 1996), and these tend to be more radio-luminous than the average. However, clear dynamical evidence exists for a relatively strong thermal wind arising from the central part of the accretion disk (Rodriguez et al 2002, Zamanov et al. 2002).

In the gas-rich and cloudy circumnuclear environments, light and low-power radio jets are readily disrupted and suffer entrainment from the surrounding material, and molecular clouds are crushed in the high-pressure environment. This is clearly demonstrated by the 2-d hydrodynamic simulations described above. However, shock velocities in Seyferts will generally be much lower than in GPS sources, and although shocks may be still important in shaping the circum-nuclear medium, photoionisation by the central nucleus appears to be much more important for its excitation.

Because clouds lying in the path of the jet and its surrounding high-pressure cocoon are crushed at relatively low velocity, any dust mixed with the cloud gas is likely to survive. Assuming that the central source produces UV photons with high local ionisation parameter, the dusty ionised gas is compressed, raising the pressure close to the ionisation front to match the radiation pressure in the EUV radiation

field Dopita et al. (2002)). This regulates the apparent ionisation parameter, and ensures that the density of the photoionized clouds varies as R^{-2} . Each photoablating cloud is surrounded by a coronal medium in which the local ionization parameter reflects the “true” ionisation parameter delivered by the central source, and each has a dusty photo-accelerated radial tail, which may be accelerated to very high velocities ($> 1000 \text{ km s}^{-1}$) (c.f. Cecil et al. 2002).

3. CONCLUSIONS

In this paper we have presented observational and theoretical evidence to support the following model for the gross features of the temporal evolution for the NLR:

1. Early on, the radio jet (if there is one), or else a strong outflow of thermal gas from the inner accretion disk pushes a strong shock into the surrounding galactic ISM.
2. Dense molecular clouds in the shocked region are crushed by radiative shocks, but along the jet axis, such clouds may be rapidly ablated and shredded by non-radiative shocks. In this phase, the optical/UV spectral signature is one of fast radiative shocks.
3. Eventually, the jet escapes from the dense region of the galaxy, and the pressure driving the shocked cocoon drains away. The wall shocks then become first radiative, and finally, momentum conserving. Shock-induced star formation may eventually occur within them, provided they become unstable to their self-gravity. The opening angle of the shocked cocoon provides the “ionization cone”. This opening angle tends to increase with time, allowing an evolution from a Seyfert II-like appearance early on, to predominantly Seyfert I-like or BLR-like spectra at late times.
4. In this phase, the crushed clouds left behind in the cocoon are photoablated by the strong UV field from the central engine, and the ablated ionized gas may be radiatively accelerated in the radial direction. In this phase the NLR presents the signatures of a photoionized plasma, and in Seyfert galaxies the coronal lines may be strong.

M. Dopita acknowledges the support of the ANU and the Australian Research Council through his ARC Australian Federation Fellowship, and both he and Ralph Sutherland acknowledge support through their ARC Discovery project DP0208445. Research by Bicknell, Saxton and Sutherland was supported by an ARC Large Grant A69905341.

REFERENCES

- Allen, M. G., Dopita, M. A., Tsvetanov, Z. I. & Sutherland, R. S. 1999, *ApJ*, 511, 686
- Axon, D. J. et al. 1998, *ApJ*, 496, L75
- Baldwin, J. A., Phillips, M. M. & Terlevich, R. 1981, *PASP*, 93, 5
- Baum, S. A., Zirbel, E. A. & O'Dea, C. P. 1995, *ApJ*, 389, 208
- Begelman, M. C. 1996, in *Cygnus A: A Study of a Radio Galaxy*, ed. C. L. Carilli & D. A. Harris (CUP: Cambridge), p209
- Best, P. N., Röttgering, H. J. A. & Longair, M. S. 2000a, *MNRAS*, 311, 1
- Best, P. N., Röttgering, H. J. A. & Longair, M. S. 2000a, *MNRAS*, 311, 23
- Bicknell, G. V., Dopita, M. A. & O'Dea, C. P. 1997, *ApJ*, 485, 112
- Bicknell, G. V., et al. 2000, *ApJ*, 540, 678
- Bicknell, G. V., Saxton, G. V., & Sutherland, R. S. 2002, *ProcASA*, (in press)
- Binette, L., Wilson, A.S. & Storchi-Bergmann, T. 1996, *A&A*, 312, 365
- Bower, G. A. et al. 1995, *ApJ*, 454, 106
- Capetti, A. et al. 1995, *ApJ*, 448, 600
- Carvalho, J. C. & O'Dea, C. P. 2002a, *ApJS*, 141, 337
- Carvalho, J. C. & O'Dea, C. P. 2002b, *ApJS*, 141, 371
- Cecil, G. et al. 2002, *ApJ*, 568, 627
- Véron, Gonçalves, & Véron-Cetty, M.-P. 1999, *A&AS*, 135, 437
- Condon, J. J.; Anderson, M. L. & Helou, G. 1991, *ApJ*, 376, 95
- de Vries, W. H. et al. 1997, *ApJS*, 110, 191
- de Vries, W. H., O'Dea, C. P., Baum, S. A., & Barthel, P. D. 1999, *ApJ*, 526, 27
- De Breuck, C. 2000, Thesis, University of Leiden.
- Dopita, M. A. & Sutherland, R. S. 1995, *ApJ*, 455, 468
- Dopita, M. A. & Sutherland, R. S. 1996, *ApJS*, 102, 161
- Dopita, M. A., Groves, B., Sutherland, R. S. & Binette, L. 2002, *ApJ*, 572, 753
- Drake, C. L., McGregor, P. J., Bicknell, G. V., & Dopita, M. A. 2002, *ProcASA* (in press).
- Evans, I. et al. 1999, *ApJ*, 521, 531
- Falcke, H., Wilson, A. S., & Simpson, C. 1998, *ApJ*, 502, 199
- Fanti, R. et al. 1990, *A&A*, 231, 333
- Francis, P. J. et al. 1991, *ApJ*, 373, 465
- Gelderman, R., & Whittle, M. 1994, *ApJS*, 91, 491
- Goldader, J. D., Joseph, R. D., Doyon, R., & Sanders, D. B. 1995, *ApJ*, 444, 97
- Haniff, C. A., Wilson, A. S., & Ward, M. J. 1988, *ApJ*, 334, 104
- Ho, L. M., Filippenko, A. V. & Sargent, W. L. 1995, *ApJS*, 98, 477
- Inskip, K. J. et al. 2002, *MNRAS* (in press), *astro-ph/0208548*.
- Inskip, K. J. et al. 2002, *MNRAS* (in press), *astro-ph/0208549*.
- Irwin, J. A., & Sarazin, C. L. 1996, *ApJ*, 471, 683.
- Kewley, L. J., Heisler, C. A., Dopita, M. A. & Lumsden, S. 2001 *ApJS*, 132, 37
- Kim, D.-C. et al. 1995, *ApJS*, 98, 129
- Kim, D.-C., Veilleux, S., Sanders, D. B. 1998, *ApJ*, 508, 627
- Lonsdale, C. J., Smith, H. J. & Lonsdale, C. J. 1993, *ApJ*, 405, 9
- O'Dea, C.P. 1998, *PASP*, 110, 493
- Osterbrock, D. E., Tran, H. D. & Veilleux, S. 1992, *ApJ*, 389, 196
- Pedlar, A. et al. 1989, *MNRAS*, 238, 863
- Phillips, M. M. et al. 1986, *AJ*, 91, 1062
- Phillips, R.B. & Mutel, R.L. 1982, *A&A*, 106, 21
- Rodriguez-Ardila, A., Viegas, S.M., Pastoriza, M.G., & Prato, L. 2002, *ApJ*, 579, 214
- Reuland, M. et al. 2003, *ApJ*, in press.
- Sanders, D. B., Soifer, B. T., Elias, J. H., Madore, B. F., Matthews, K., Neugebauer, G. & Scoville, N. Z. 1988, *ApJ* 325, 74
- Scheuer, P. A. G. 1982, in *IAU Symp.97: Extragalactic Radio Sources*, ed D. S. Heeschen & C. M. Wade (Dordrecht:Reidel), p163
- Steffen, W., Gomez, J. L., Williams, R. J. R., Raga, A. C., & Pedlar, A. 1997, *MNRAS*, 286, 1032
- Sutherland, R. S. Bisset, D. K., & Bicknell, G. V. 2002, *ApJ*, submitted.
- Veilleux, S. & Osterbrock, D. E. 1987, *ApJS*, 63, 295
- Veilleux, S. 1991a *ApJS*, 75, 357,
- Veilleux, S. 1991b, *ApJS*, 75, 383
- Veilleux, S. 1991c *ApJ*, 369, 331
- Veilleux, S., Kim, D.-C. & Sanders, D. B. 1999, *ApJ*, 522, 113
- Véron, P., Gonçalves, A. C. & Véron-Cetty, M.-P. 1997, *A&A*, 319, 52
- Véron, P. & Véron-Cetty, M.-P. 2000, *A&ARev.*, 10, 81
- Véron, P. & Véron-Cetty, M.-P. 1986, *A&A*, 161, 145
- Whittle, M. et al. 1988, *ApJ*, 326, 125
- Whittle, M. 1996, *ApJS*, 79, 49
- Wilkinson, P.N. et al. 1994, *ApJ*, 432, L87
- Zamanov, R. et al. 2002, *ApJ*, 576, L9

M. A. Dopita, G. V. Bicknell, R. S. Sutherland & C. J. Saxton: Research School of Astronomy & Astrophysics, Institute of Advanced Studies, The Australian National University, Cotter Road, Weston Creek, ACT 2611, Australia (Michael.Dopita@anu.edu.au).